

Observation of a 1750 MeV/ c^2 Enhancement in the Diffractive Photoproduction of K^+K^-

The FOCUS Collaboration[★]

J. M. Link^a M. Reyes^a P. M. Yager^a J. C. Anjos^b I. Bediaga^b
 C. Göbel^b J. Magnin^b A. Massafferri^b J. M. de Miranda^b
 I. M. Pepe^b A. C. dos Reis^b S. Carrillo^c E. Casimiro^c
 E. Cuautle^c A. Sánchez-Hernández^c C. Uribe^c F. Vázquez^c
 L. Agostino^d L. Cinquini^d J. P. Cumalat^d B. O'Reilly^d
 J. E. Ramirez^d I. Segoni^d J. N. Butler^e H. W. K. Cheung^e
 G. Chiodini^e I. Gaines^e P. H. Garbincius^e L. A. Garren^e
 E. Gottschalk^e P. H. Kasper^e A. E. Kreymer^e R. Kutsche^e
 L. Benussi^f S. Bianco^f F. L. Fabbri^f A. Zallo^f C. Cawfield^g
 D. Y. Kim^g K. S. Park^g A. Rahimi^g J. Wiss^g R. Gardner^h
 A. Kryemadhi^h C. H. Changⁱ Y. S. Chungⁱ J. S. Kangⁱ
 B. R. Koⁱ J. W. Kwakⁱ K. B. Leeⁱ K. Cho^j H. Park^j
 G. Alimonti^k S. Barberis^k A. Cerutti^k M. Boschini^k
 P. D'Angelo^k M. DiCorato^k P. Dini^k L. Edera^k S. Erba^k
 M. Giammarchi^k P. Inzani^k F. Leveraro^k S. Malvezzi^k
 D. Menasce^k M. Mezzadri^k L. Moroni^k D. Pedrini^k
 C. Pontoglio^k F. Prelz^k M. Rovere^k S. Sala^k
 T. F. Davenport III^ℓ V. Arena^m G. Boca^m G. Bonomi^m
 G. Gianini^m G. Liguori^m M. M. Merlo^m D. Pantea^m
 S. P. Ratti^m C. Riccardi^m P. Vitulo^m H. Hernandezⁿ
 A. M. Lopezⁿ H. Mendezⁿ L. Mendezⁿ E. Montielⁿ D. Olayaⁿ
 A. Parisⁿ J. Quinonesⁿ C. Riveraⁿ W. Xiongⁿ Y. Zhangⁿ
 J. R. Wilson^o T. Handler^p R. Mitchell^p D. Engh^q M. Hosack^q
 W. E. Johns^q M. Nehring^q P. D. Sheldon^q K. Stenson^q
 E. W. Vaandering^q M. Webster^q M. Sheaff^r

^aUniversity of California, Davis, CA 95616

^bCentro Brasileiro de Pesquisas Físicas, Rio de Janeiro, RJ, Brasil

^cCINVESTAV, 07000 México City, DF, Mexico

[★] see <http://www-focus.fnal.gov/authors.html> for additional author information.

- ^d*University of Colorado, Boulder, CO 80309*
^e*Fermi National Accelerator Laboratory, Batavia, IL 60510*
^f*Laboratori Nazionali di Frascati dell'INFN, Frascati, Italy I-00044*
^g*University of Illinois, Urbana-Champaign, IL 61801*
^h*Indiana University, Bloomington, IN 47405*
ⁱ*Korea University, Seoul, Korea 136-701*
^j*Kyungpook National University, Taegu, Korea 702-701*
^k*INFN and University of Milano, Milano, Italy*
^l*University of North Carolina, Asheville, NC 28804*
^m*Dipartimento di Fisica Nucleare e Teorica and INFN, Pavia, Italy*
ⁿ*University of Puerto Rico, Mayaguez, PR 00681*
^o*University of South Carolina, Columbia, SC 29208*
^p*University of Tennessee, Knoxville, TN 37996*
^q*Vanderbilt University, Nashville, TN 37235*
^r*University of Wisconsin, Madison, WI 53706*

Abstract

Using the FOCUS spectrometer with photon beam energies between 20 and 160 GeV, we confirm the existence of a diffractively photoproduced enhancement in K^+K^- at $1750 \text{ MeV}/c^2$ with nearly 100 times the statistics of previous experiments. Assuming this enhancement to be a single resonance with a Breit-Wigner mass shape, we determine its mass to be $1753.5 \pm 1.5 \pm 2.3 \text{ MeV}/c^2$ and its width to be $122.2 \pm 6.2 \pm 8.0 \text{ MeV}/c^2$. We find no corresponding enhancement at $1750 \text{ MeV}/c^2$ in K^*K , and again neglecting any possible interference effects we place limits on the ratio $\Gamma(X(1750) \rightarrow K^*K)/\Gamma(X(1750) \rightarrow K^+K^-)$. Our results are consistent with previous photoproduction experiments, but, because of the much greater statistics, challenge the common interpretation of this enhancement as the $\phi(1680)$ seen in e^+e^- annihilation experiments.

PACS numbers: 13.25 13.60.L 14.40

1 Motivation

Previous photoproduction experiments have consistently observed an enhancement in K^+K^- at a mass near $1750 \text{ MeV}/c^2$ (referred to here as the $X(1750)$) [1–3]. However, with signals consisting of only around 100 events and with a statistical significance of the enhancement of, at best, only 3.5σ over background, these experiments have suffered from a lack of statistics. Due to the large statistical errors on the mass of the enhancement, and assuming that

a diffractively photoproduced meson is a vector, this enhancement has been identified with the $\phi(1680)$ seen in e^+e^- annihilation [4–7], which is a candidate radial excitation of the $\phi(1020)$. The present analysis challenges this interpretation of the 1750 MeV/ c^2 signal on two grounds.

First, assuming the enhancement is a single resonance, we determine a mass of $1753.5 \pm 1.5 \pm 2.3$ MeV/ c^2 from our sample of more than 10,000 signal events, which is clearly inconsistent with 1680 MeV/ c^2 . One previous photoproduction experiment [1], using less than 50 signal events and guided by e^+e^- annihilation results, presented a mass of 1690 ± 10 MeV/ c^2 after an analysis based on a model including interference and Deck-like effects. With orders of magnitude more statistics, we are unable to reproduce this result.

Second, although e^+e^- annihilation experiments report the dominant decay mode of the $\phi(1680)$ to be K^*K [7], as expected theoretically for the radial excitation of the $\phi(1020)$ [8–10], we find no evidence for a photoproduced enhancement in K^*K corresponding to the photoproduced enhancement in K^+K^- . We put a low upper limit on the ratio $\Gamma(X(1750) \rightarrow K^*K)/\Gamma(X(1750) \rightarrow K^+K^-)$.

2 Data Sample

The data for this analysis was collected by the Wideband photoproduction experiment FOCUS during the Fermilab 1996–1997 fixed-target run. FOCUS is a considerably upgraded version of the previous E687 photoproduction experiment [11]. A forward multi-particle spectrometer is used to measure the interactions of high energy photons on a segmented BeO target.

The FOCUS detector is a large aperture, fixed-target spectrometer with excellent vertexing, particle identification [12], and reconstruction capabilities for photons, π^0 's, and K_S [13]. A photon beam is derived from the bremsstrahlung of secondary electrons and positrons with an approximately 300 GeV end-point energy produced from the 800 GeV Tevatron proton beam. The charged particles which emerge from the target are tracked by two systems of silicon microvertex detectors. The upstream system, consisting of 4 planes (two views in 2 stations), is interleaved with the experimental target, while the other system lies downstream of the target and consists of twelve planes of microstrips arranged in three views. The momentum of a charged particle is determined by measuring its deflections in two analysis magnets of opposite polarity with five stations of multiwire proportional chambers. Three multicell threshold Cerenkov counters are used to discriminate between pions, kaons, and protons.




Fig. 1. The K^+K^- sample with no cut on p_T .

The K^+K^- and $K_S K^\pm \pi^\mp$ data samples used in this paper require a single vertex in the target, no electromagnetic energy apart from that associated with the reconstructed tracks, and require all tracks to be singly linked between the upstream and downstream tracking systems. Events with additional reconstructed tracks are rejected, making the data samples as exclusive as possible. The Cerenkov identification of the kaons in both diffractive final states limits the photon energy range to ≤ 160 GeV.

3 Mass and Width Measurement

Our sample of K^+K^- events, selected using the criteria described above, shows a clear $\phi(1020)$ signal dominating the spectrum (Figure 1). The diffractive component of the production of the $\phi(1020)$ shows up as a peak in the p_T spectrum (Figure 2.a). Cutting around this peak by requiring $p_T < 0.15$ GeV/c, we select a diffractive sample of K^+K^- events, in which a clear enhancement appears in the mass spectrum near 1750 MeV/c² (Figure 3.a). Figure 3.b confirms that the enhancement appears at only low p_T . Plotting the p_T spectra in the 1750 region (1640–1860 MeV/c²) and in the two sideband regions (1500–1600 MeV/c² and 1900–2100 MeV/c²), it is seen that the 1750 region has a peak in the p_T spectrum in nearly the same place as the $\phi(1020)$ peak, but the sideband regions have significantly smaller p_T peaks (Figure 2.a), indicating that the background under the X(1750) signal is largely non-diffractive.

The t' spectrum ($t' \equiv |t| - |t|_{min} \approx p_T^2$ and $t \equiv (P_\gamma - P_{KK})^2$, where P_γ and P_{KK} are the four-momenta of the photon beam and the KK system, respectively) for K^+K^- masses between 1640 and 1860 MeV/c² has been fit with two exponentials (Figure 2.b). The diffractive exponential for this region has a slope of 69.2 ± 2.1 GeV⁻²c² and the background exponential has a slope of




Fig. 2. (a) The K^+K^- p_T spectra. The solid line is the p_T spectrum for the $\phi(1020)$. The top dotted line is the p_T spectrum for K^+K^- masses between 1640 and 1860 MeV/ c^2 ; the middle is the left sideband (1500–1600 MeV/ c^2); and the bottom is the right sideband (1900–2100 MeV/ c^2). (b) The t' spectrum for K^+K^- masses between 1640 and 1860 MeV/ c^2 fitted using two exponentials, $A_1e^{-b_1t'} + A_2e^{-b_2t'}$. In each plot, the vertical line represents the p_T cut used in this analysis.

$4.17 \pm 0.21 \text{ GeV}^{-2}c^2$. For comparison, the $\phi(1020)$ signal has a diffractive slope of $77.71 \pm 0.59 \text{ GeV}^{-2}c^2$ and a background slope of $1.71 \pm 0.14 \text{ GeV}^{-2}c^2$. The steep diffractive slopes are characteristic of exclusive diffractive photoproduction off of nuclear targets.

Fitting the 1750 MeV/ c^2 mass region with a non-relativistic Breit-Wigner distribution and a quadratic background, we find

$$\text{Yield} = 11,700 \pm 480 \text{ Events}$$

$$M = 1753.5 \pm 1.5 \pm 2.3 \text{ MeV}/c^2$$

$$\Gamma = 122.2 \pm 6.2 \pm 8.0 \text{ MeV}/c^2$$

Because the acceptance of the detector is flat in this region, as determined




Fig. 3. (a) The K^+K^- mass spectrum with the requirement that $p_T < 0.15 \text{ GeV}/c$. The spectrum is fit with a non-relativistic Breit-Wigner distribution and a quadratic background. The dotted line is the Monte Carlo efficiency on a scale from 0 to 100%. (b) The solid line is the K^+K^- mass spectrum with the requirement that $p_T < 0.15 \text{ GeV}/c$. The dotted line is the K^+K^- mass spectrum with $p_T > 0.15 \text{ GeV}/c$ scaled to the size of the low p_T spectrum for comparison. (c) The data and fit after subtracting the quadratic polynomial background shape. (d) The data minus the fit.

by a full Monte Carlo simulation, the fit was performed on the uncorrected mass spectrum. Further, since Monte Carlo studies of the detector have shown that the K^+K^- mass resolution in the $1750 \text{ MeV}/c^2$ region is better than $10 \text{ MeV}/c^2$, which is much less than the width of the $X(1750)$, resolution effects have been neglected in the fit. (Similar Monte Carlo studies have accurately predicted the D^0 mass resolution in the decay $D^0 \rightarrow K^+K^-$, and find a compatible resolution [14].) The systematic errors were determined by varying the p_T cut, the Cerenkov cuts, the form of the Breit-Wigner shape (non-relativistic, and relativistic $L = 0, 1, 2$), and the form of the background shape, and include the systematic uncertainty in our mass scale.

There is a region near $1600 \text{ MeV}/c^2$ where there is some discrepancy in our fit to the K^+K^- mass spectrum. The residuals show that the statistical significance of this discrepancy is not strong (Figure 3.d). It has been found that several different interference scenarios can improve the fit. These include interference with the K^+K^- continuum and interference between the $X(1750)$




Fig. 4. (a) The $K_S K \pi$ sample with no cut on p_T . (b) The $K_S K \pi$ sample with $p_T < 0.15$ GeV/c. (c) The $K_S \pi^\pm$ mass from (b). (d) The $K^\pm \pi^\mp$ mass from (b).

and a second resonance with lower mass. The goodness of the fits, however, does not allow us to discriminate between solutions, and we find no physics motivation for picking one solution over another. In all scenarios, the mass of the X(1750) exceeds 1747 MeV/c². If the X(1750) has $J^{PC} = 1^{--}$ and if the $\phi(1680)$ were also photoproduced, then we would expect some distortion of the $K^+ K^-$ mass shape. However, the apparent absence of the $\phi(1680)$ in its dominant $K^* K$ mode and the measured ratio of branching fractions [7] limit the amplitude in the $K^+ K^-$ state so that the distortion of the expected line shape is negligible. Our mass and width measurements and our determination of systematic errors assume the production of a single, non-interfering resonance.

4 Branching Ratio

Figure 4 shows our $K_S K \pi$ sample and the two K^* combinations. The $K_S K^\pm \pi^\mp$ spectrum shows the classic “D” and “E” regions [15].

After requiring $p_T < 0.15$ GeV/c (the same p_T cut imposed on the $K^+ K^-$ sample), and requiring a K^* , the two distinct $K^* K$ spectra were fit individually (Figure 5). There is no obvious X(1750) signal in either of the two $K^* K$




Fig. 5. Fits to K^*K using a Breit-Wigner distribution for the X(1750) region with mass and width fixed from the fit to K^+K^- , a second non-interfering Breit-Wigner distribution around 1630 MeV/c², and a quadratic background. The dotted lines are the Monte Carlo efficiencies on a scale from 0 to 20%. (a) $K^{*0}K_S$ with K^{*0} to $K^\pm\pi^\mp$. (b) $K^{*\pm}K^\mp$ with $K^{*\pm}$ to $K_S\pi^\pm$.

modes. In order to place upper limits on the K^*K/K^+K^- branching ratios of the X(1750), an estimate of the background is needed. The presence of a slight enhancement somewhat below the $\phi(1680)$ region introduces some ambiguity in estimating the background in the X(1750) region. In order to make a conservative estimate of the background, we have used a fit which includes a second, unconstrained, non-interfering resonance in the 1630 MeV/c² region as well as the X(1750) with mass and width fixed from the fit to the K^+K^- mode. The size of the X(1750) resonant component was unconstrained. These fits provide an estimate of the number of events above background in the X(1750) region, -123 ± 120 events in the $K^{*0}K_S$ mode and 106 ± 117 in the $K^{*\pm}K^\mp$ mode.

The efficiencies of the K^+K^- and K^*K final states were determined by Monte Carlo simulations in bins of p_T , beam energy, and mass. As the spin of the X(1750) is still uncertain, several decay angular distributions were simulated. Using the highest efficiency for K^+K^- (Figure 3.a) and the lowest for K^*K (Figure 5) and correcting for the K_S unseen decay mode, we have found an upper limit on the following relative branching ratios

$$\frac{\Gamma(X(1750) \rightarrow \bar{K}^{*0}K^0 \rightarrow K^-\pi^+K_S + c.c.)}{\Gamma(X(1750) \rightarrow K^+K^-)} < 0.065 \text{ at } 90\% \text{ C.L.}$$

$$\frac{\Gamma(X(1750) \rightarrow K^{*+}K^- \rightarrow K_S\pi^+K^- + c.c.)}{\Gamma(X(1750) \rightarrow K^+K^-)} < 0.183 \text{ at } 90\% \text{ C.L.}$$

The confidence limits were set using the Feldman-Cousins methodology [16]. The two relative branching ratios were measured to be -0.083 ± 0.081 and

0.065 ± 0.072 , respectively.

5 Discussion

Because of the large discrepancies in mass and relative branching fractions to K^+K^- and K^*K , we do not believe it is reasonable to identify the X(1750) with the $\phi(1680)$. In fact, because the mass of the X(1750) is significantly higher than all known vector mesons, the most massive of which are the $\omega(1650)$, $\phi(1680)$, and $\rho(1700)$, an interpretation claiming the X(1750) is some combination of interfering vector mesons also seems highly unlikely. The interpretation of the X(1750) remains uncertain.

6 Acknowledgements

We wish to acknowledge P. Page and T. Barnes for valuable discussions, the assistance of the staffs of Fermi National Accelerator Laboratory, the INFN of Italy, and the physics departments of the collaborating institutions. This research was supported in part by the U. S. National Science Foundation, the U. S. Department of Energy, the Italian Istituto Nazionale di Fisica Nucleare and Ministero dell'Università e della Ricerca Scientifica e Tecnologica, the Brazilian Conselho Nacional de Desenvolvimento Científico e Tecnológico, CONACyT-México, the Korean Ministry of Education, and the Korean Science and Engineering Foundation.

References

- [1] OMEGA Collaboration, D. Aston et al., Phys. Lett. B 104 (1981) 231.
- [2] OMEGA Collaboration, M. Atkinson et al., Zeit. Phys. C 27 (1985) 233.
- [3] FNAL E401 Collaboration, J. Busenitz et al., Phys. Rev. D 40 (1989) 1.
- [4] D.E. Groom et al. (Particle Data Group), Eur. Phys. J. C 15, (2000) 1.
- [5] A.B. Clegg, A. Donnachie, Zeit. Phys. C 62 (1994) 455.
- [6] DM2 Collaboration, D. Bisello et al., Zeit. Phys. C 52 (1991) 227.
- [7] DM1 Collaboration, J. Buon et al., Phys. Lett. B 118 (1982) 221.

- [8] S. Godfrey and N. Isgur, Phys. Rev. D 32 (1985) 189.
- [9] H.G. Blundell and S. Godfrey, Phys. Rev. D 53 (1996) 3700.
- [10] H.G. Blundell, S. Godfrey, and B. Phelps, Phys. Rev. D 53 (1996) 3712.
- [11] FNAL E687 Collaboration, P.L. Fabretti et al., Nucl. Instrum. Methods A 320 (1992) 519.
- [12] FOCUS Collaboration, J.M. Link et al., Nucl. Instrum. Methods A 484 (2002) 270.
- [13] FOCUS Collaboration, J.M. Link et al., Nucl. Instrum. Methods A 484 (2002) 174.
- [14] FOCUS Collaboration, J.M. Link et al., Phys. Lett. B 491 (2000) 232.
- [15] S. Godfrey and J. Napolitano, Rev. Mod. Phys. 71 (1999) 1411.
- [16] G.J. Feldman and R.D. Cousins, Phys. Rev. D 57 (1998) 3873.